















Principal Investigators:
Frank Muller-Karger (USF)
Maria Kavanaugh, Robert Cowen (OSU)
Francisco Chavez (MBARI) Robert Miller (UCSB)
Jeffrey Runge (UMaine)
Katrin Iken (UAF)
Enrique Montes (USF)
Emmett Duffy (SI)


































Huge investments have gone into ocean observing systems — But there is still no systematic effort to observe life in the sea

MBON Goal: Enable the effort to characterize how marine biodiversity is changing and how it affects us

Essential Variables



EBVs are not measured. They are derived Scenarios for biodiversity & ecosystem services (e.g. for IPBES) (multidimensional data 'cubes') High-level indicators of biodiversity Ancillary attributes Ecosystem-service valuation & other data Observations Observations of policy of drivers & & management **Essential Biodiversity** pressures Variables Genetic composition Species populations Species traits Ecosystem function Primary observations of


Pereira et al. (2013) GEO BON EBVs

Framework for Ocean Observing (2012) Global Ocean Observing System (GOOS) Essential Ocean Variables (EOVs) EOVs are the primary observations needed to model and build the EBVs (multidimensional data cubes)




Biology and Ecosystem Essential Ocean Variables (EOVs)

FUNCTIONAL GROUPS Phytoplankton Fish Microbes Zooplankton Benthic invertebrates Turtle-Bird-Mammal **Distribution and Abundance** Distribution and Abundance **Diversity and Biomass Diversity and Biomass** Distribution and Abundance **Diversity and Biomass HABITAT STATE Hard Coral** Macroalgae Seagrass Mangrove **Cover and Composition** Cover and composition Cover and composition Cover and composition



http://marinebon.org

Satellite-derived Chlorophyll distribution



http://marinebon.org





Satellite-derived Chlorophyll distribution

2021 United Nations Decade of Ocean Science for Sustainable Development

Evolving technology matrix

	Microbes/ Phyto	Zooplankton	Fish	Top Predators	Benthos, habitat forming
Optics/Imaging	X	X	X Benthic		X
Animal tracking (satellite, underwater)			X	X	
Acoustics		X active	X Active, passive	X Tags, passive	X Active, passive (noise)
Genomics	X	X	X	X	X
Platforms with samplers	AUVs, floats, moorings, satellites	AUVs, moorings	AUVs, moorings	AUVs, moorings, tags	AUVs, moorings, satellites
Data and visualization	X	X	X	X	X


OCEANOGRAPHY SOCIETY VOL.34, NO.2, JUNE 2021




MBON

MARINE BIODIVERSITY OBSERVATION NETWORK

AN OBSERVING SYSTEM FOR LIFE IN THE SEA






Southern California Bight MBON

Central California MBON


Northern California Current MBON



Santora et al., 2021

https://doi.org/10.5670/oceanog.2021.212

Kavanaugh et al., 2021



Finding:

- Record high diversity in 2014–2016
- N Pacific heatwave compressed upwelling habitat against the coast

















Finding:

Borealization of the Chukchi Sea Shelf


Northward expansion of predators:

- demersal fish and large invertebrates
- reorganization of benthic food web

Arctic Marine Biodiversity Observing Network (AMBON)




Mueter et al., 2021 https://doi.org/10.5670/oceanog.2021.213



Bottom-up drivers in the Chukchi Sea (green arrows) 10

Florida Florida Keys National Marine Sanctuary NATIONAL MARINE SANCTUARIES **No-Take Marine Zones** FKNMS Grecian $N (\pm SE)$ Hens and Chickens Boundary of FKNMS Sanctuary Preservation Area ☐ Research Only (RO) Newfound Western Sambo (ER) Harbor $g/m^2 (\pm SE)$ East Dry 81°0'0"W 82°0'0"W

South Florida MBON




Medina et al., 2021 https://doi.org/10.5670/oceanog.2021.214


Finding:

Diversity difference between protected and unprotected areas has decreased over time

High relief reefs have higher diversity







1.65

2000


2004

2008

Year


2012

2016



Gulf of Maine MBON

Warming of the Gulf of Maine (GoM) affects valuable fisheries and marine mammals (critically endangered N Atlantic right whales)




The *Calanus* Index. based on monthly measurements of *C. finmarchicus abundance*, W Gulf

Abundance levels now 30-50% of historical levels in late summer through winter. [y-axis is log scale](Ji et al. 2021 and GoM MBON data).

Distribution of 16 groups of similar coastlines

Global Efforts



Sayre et al., 2021 https://doi.org/10.5670/oceanog.2021.219



VARIABLE	MEAN	MEDIAN	STANDARD DEVIATION	RANGE	PREDOMINANT CLASS (% OF SEGMENTS)
Slope (%)	8.84	4.52	11.25	100.08	Flat (71)
Sinuosity (unitless)	10.66	1.41	98.18	1212.32	Straight (53)
Erodibility	NA	NA	NA	NA	Low (94)
Temperature and Moisture Regime	NA	NA	NA	NA	Tropical Moist (94)
River Discharge (unitless)	0.0002	0.00002	0.0005	0.002	Moderate River Discharge (70)
Wave Height (m)	0.41	0.41	0.16	0.48	Low Wave Energy (68)
Tidal Range (m)	1.00	1.02	0.25	1.21	Moderately Tidal (52)
Marine Physical Environment	NA	NA	NA	NA	Euhaline-Oxic-Warm (100)
Turbidity (m ⁻¹)	0.27	0.19	0.20	0.53	Clear (41)
Chlorophyll (µg L ⁻¹)	2.30	2.28	0.45	3.71	Moderate Chlorophyll (69)

Coastline characteristics of the Florida Keys National Marine Sanctuary (Florida Keys US MBON site)

Output:

CSU - Coastal Segment Units

Coastal and Marine Ecological Classification Standard (CMECS) labels for global coastal segments at 1 km

Output:

MBON and OBIS/GBIF collaboration


Mapping data flow: collection to publication

Examples:

- Rockfish (California/Pacific)
- Reef fish (Florida keys/Atlantic)
- Satellite data (early alert dashboards)
- Pole to Pole Data Viewer
- IOOS MBON Portal

DATA VIEW Global **RAW DATA CLEAN & DATA SERVER Explorer TYPE STANDARDIZED** DATA **Infographics OBIS** Taxa Darwin **Occurrences** Core Pole to Pole Website **ERDDAP Early Alerts Dashboards** Satellite Climate & Data **Forecast Portal MBONIOOS Data Views Portal**


Data Management and Interactive Visualizations




Benson et al., 2021 https://doi.org/10.5670/oceanog.2021.220



Addressing Sanctuary Needs: data tools





Pole to Pole MBON

Developing Essential Biodiversity Variables (EBVs)
















100°W 75°W 50°W 25°W O Collaborations/monitoring sites

"Ecosystem Extent" EBV using satellite-derived seascape maps Capacity Building – Field and satellite data / applications





Challenge 2:
Develop solutions
to ecosystem stressors



Challenge 9: Develop capacity and equitable access



Challenge 10:

Understand values and services, humanity's relationship with ocean



Our Partners

























And a long list of academic and government partners

https://marinelife203

Thank You!

Frank Muller-Karger carib@usf.edu

http://marinebon.org

#MarineLife2030

https://marinelife2030.org/



